

Longitudinal Distribution of Fishes in the Sawmill River, New York

Study Area:

The Sawmill River is a tributary of the Hudson River, in Westchester County, New York. It flows for approximately 22 river miles, from its' source in the Town of New Castle, to the Hudson River in the City of Yonkers. The mouth of the Sawmill River is tidal and receives an influx of water from the Hudson River. It displays a distinct dendritic drainage pattern in its' upper course and a long narrow drainage basin in its' southern reaches. The Sawmill River Parkway runs along the bank of the river and criss-crosses it at several places along its' length. The northern reaches of the Sawmill River course through densely populated suburban development, eventually reaching the City of Yonkers before flowing into the Hudson River. The Sawmill River is widest, at its' confluence with the Hudson River. Upstream areas are 3-8 meters wide with depth less than 1 meter. The stream follows a typical pattern of alternating pools and riffles. The bottom varies from mud and sand to cobble and boulder.

Materials and Methods:

Ten sites were selected for sampling, based on ease of access from the highway (fig. 1). Station 1 was at the confluence of the Sawmill and Hudson Rivers while Station 10 was located at headwaters of the Sawmill (Town of New Castle).

Each station was sampled using a 1 x 5.5 meter seine, with the exception of Dock Street, which was sampled using a fyke net. The fyke net was placed in position at low tide and retrieved on the following low tide. Sampling was initiated in April 1989 and terminated in August 1989. All specimens collected, were fixed in 10% formalin and later transferred to 70% ethanol. Specimens were identified and counted. Other data such as length, weight, sex and food habits are presently being collected for use in later studies.

Data was quantified, in part, using a graphical method to assess faunal dominance (Guillory, 1977), which is modified from Saunders (1960) and Ono (1960). The ichthyofauna was characterized by plotting an expression of frequency of occurrence vs. an expression of percent abundance. The fish were categorized as abundant, common,

rare or Hudson River.

Results and Discussion:

A total of 1967 specimens were collected representing 12 families and 22 species (Table 1). Figure 1 shows that the most speciose station was Dock Street (Station 1). Fourteen different species were collected at this one site. Of these 14 species, 8 were found only at Dock Street, the other 6 were found at one or more stations along the length of the river. These data show that Hudson River fish utilize the mouth of the Sawmill River. Further, longterm studies are needed however to determine for what purpose these fish utilize the mouth of the Sawmill River. Among the possibilities are feeding, breeding or nursery areas.

Although the Sawmill River appears to be a rather diverse system, there are species, quite common in other Westchester County streams, that may have been eliminated from the Sawmill River i.e., fall fish (Semotilus corporalis) and the cutlips minnow (Exoglossum maxillingua). Schmidt and Samaritan (1984) found a similar situation in the Bronx River. They attributed the loss of these species to poor water quality. One introduced species, the European

bitterling, Rhodeus sericeus, has also disappeared from the Sawmill River, although it is still present in the Bronx River (Schmidt et al, 1981). Early records such as Greely (1936) and later collections (Cornell University 23611; 1951) still listed the species as present. Efforts to collect this species, including my own have proved to no avail.

Three species in the Sawmill River are not native to the northeastern United States: largemouth bass (Micropterus salmoides), carp (Cyprinus carpio) and the rainbow trout (Salmo gairdneri). This represents about 14% of the fish fauna of the river.

The relative abundance of calculations indicated that the blacknose dace (Rhinichthys atratulus) and the white sucker (Catostomus commersoni) were the most abundant species (Fig. 2). Both these species were collected in 8 out of 10 stations. Relative abundance as calculated by the Guillery (1977) method also listed 7 species as common, and 4 species as rare (Fig. 2). Upon examining the graph in figure 2, it became obvious that a fourth cluster of points was evident. I chose to refer to this cluster as the Hudson River group. With the exception

of one species (Esox americanus, redfin pickerel) all the species in this cluster were collected at Dock Street and represent the influence of the Hudson River, upon the Sawmill River.

References

- Greely, J.R. (1936). Fishes of the area with an annotated list. In: A biological survey of the lower Hudson watershed. Suppl. 26th Ann. Rept., NY Conserv. Dept., part II. pp. 45-85.
- Guillory, V. (1977). Graphical method to assess faunal dominance. Proc. Ann. Conf. SE Assoc. Fish and Wildlife Agencies. 31: 563-564.
- Ono, U. (1960). An ecological study of the Brachyuran community on Tomioko Bay, Amkusia, Rec. Oceanogr. Works in Japan 5: 199-210.
- Sanders, H. (1960). Benthic studies in Buzzards Bay. III. The structure of the soft-bottom community. Limn. Oceanogr. 5: 138-153.
- Schmidt, R.E. et al. (1981). Status of the bitterling, Rhodeus sericeus, in Southeastern New York. Copeia 1981(2): 481-482.
- Schmidt, R.E. and Samaritan, J. (1984). Fishes of an urban upstream; the Bronx River, New York. Northeastern Environ. Sci. 3(1): 3-7.

Figure 1. Distribution of Fishes of the Sawmill River.
1989

Table 1. Species of Fish Collected in the Sawmill River (Total Species = 22)

A. Family Cyprinidae (Minnows)

1. Longnose Dace Rhinichthys cataractae
2. Blacknose Dace Rhinichthys atratulus
3. Common Shiner Notropis cornutus
4. Golden Shiner Notemigonus crysoleucas
5. Carp Cyprinus carpio
6. Creek Chub Semotilus atromaculatus

B. Family Centrarchidae (Sunfishes and Black Basses)

1. Pumpkinseed Sunfish Lepomis gibbosus
2. Bluegill Sunfish Lepomis macrochirus
3. Redbreasted Sunfish Lepomis auitus
4. Largemouth Bass Micropterus salmoides

C. Family Moronidae (Temperate Basses)

1. White Perch Morone americana
2. Striped Bass Morone saxatilis

D. Family Clupeidae (Herrings)

1. American Shad Alosa sapidissima
2. Alewife Alosa pseudoharengus

E. Family Catostomidae (Suckers)

1. White Sucker Catostomus commersoni

Table 1. Continued:

F. Family Salmonidae (Salmon and Trout)

1. Rainbow Trout Salmo gairdneri

G. Family Percidae (Perches)

1. Tessellated Darter Etheostoma olmstedii

H. Family Gadidae (Codfishes)

1. Atlantic tomcod Microgadus tomcod

I. Family Pomatomidae (Bluefishes)

1. Bluefish Pomatomus saltatrix

J. Family Anguillidae (Eels)

1. American eel Anguilla rostrata


K. Family Sciaenidae (Drums)

1. Spot Leiostomus xanthurus

L. Family Esocidae (Pikes)

1. Redfin Pickerel Esox americanus

Figure 2. Relative Abundance of Various Species Collected on the Sawmill River
 (Abundance estimated by method of Gillory, 1977).



KEY TO FISH SPECIES

Abundant	1. Blacknose Dace	14. Tomcod
	2. White Sucker	15. Spot
	3. Common Shiner	16. White Perch
Common	4. Tessellated Darter	17. Striped Bass
	5. Golden Shiner	18. American Eel
	6. Bluegill Sunfish	19. Redfin Pickerel*
	7. Creek Chub	20. Bluefish
	8. Longnose Dace	21. Alewife
	9. Redbreasted Sunfish	22. American Shad
Rare	10. Pumpkinseed Sunfish	
	11. Largemouth Bass	
	12. Carp	
	13. Rainbow Trout	

* All fish species labeled HUDSON RIVER were collected at the confluence of the Sawmill and Hudson Rivers, with the exception of the Redfin Pickerel.